

Table 2-4
Comparison of Estimates of Water in Cyclic and Local Storage Accounts
with and without Losses to the Santa Ana River
(acre-ft)

Year	----- Cyclic and Local Storage Accounts -----					
	Put	Take	Watermaster Estimated End of Period Storage	Losses to Baseflow	Accounting for Losses Estimated End of Period Storage	Cumulative Losses
1978 / 79	16,074	0	16,074	163	15,911	-163
1979 / 80	19,898	10,678	25,295	417	24,715	-580
1980 / 81	12,665	3,021	34,938	600	33,759	-1,179
1981 / 82	6,015	2,454	38,500	721	36,599	-1,901
1982 / 83	20,345	0	58,846	949	55,995	-2,850
1983 / 84	19,158	0	78,004	1,331	73,822	-4,181
1984 / 85	25,986	615	103,375	1,756	97,437	-5,938
1985 / 86	18,192	106	121,461	2,162	113,362	-8,099
1986 / 87	31,257	14,021	138,698	2,476	128,122	-10,575
1987 / 88	58,037	17,153	179,582	3,016	165,990	-13,591
1988 / 89	43,990	31,983	191,588	3,491	174,505	-17,083
1989 / 90	26,742	34,774	183,555	3,461	163,012	-20,544
1990 / 91	34,451	5,877	212,129	3,599	187,986	-24,143
1991 / 92	83,614	66,103	229,640	3,994	201,503	-28,137
1992 / 93	30,388	23,028	237,000	4,165	204,698	-32,302
1993 / 94	32,807	21,889	247,918	4,266	211,350	-36,568
1994 / 95	30,333	6,288	271,963	4,534	230,861	-41,102
1995 / 96	38,488	34,785	275,666	4,724	229,840	-45,826
1996 / 97	20,698	22,301	274,063	4,649	223,587	-50,476